### **PCT**

# WORLD INTELLECTUAL PROPERTY ORGANIZATION International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT) (51) International Patent Classification 7: (11) International Publication Number: C07D 405/12, A61K 31/4525 A1 (43) International Publication Date: 18 May 2000 (18.05.00) (21) International Application Number: (81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, PCT/GB99/03664 (22) International Filing Date: 5 November 1999 (05.11.99) KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, (30) Priority Data: 9824298.5 5 November 1998 (05.11.98) (71) Applicant (for all designated States except US): SMITHKLINE CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, BEECHAM PLC [GB/GB]; New Horizons Court, Brentford. Middlesex TW8 9EP (GB). GN, GW, ML, MR, NE, SN, TD, TG). (72) Inventors; and (75) Inventors/Applicants (for US only): CAMBURN, Ian, David **Published** [GB/GB]; SmithKline Beecham Pharmaceuticals, Clarendon With international search report. Road, Worthing, West Sussex BN14 8QH (GB). MERRI-FIELD, David, Roy [GB/GB]; SmithKline Beecham Pharmaceuticals, New Frontiers Science Park South, Third Avenue, Harlow, Essex CM19 5AW (GB). VALDER, Christopher, Edmund [GB/GB]; SmithKline Beecham Pharmaceuticals, New Frontiers Science Park South, Third Avenue, Harlow, Essex CM19 5AW (GB). (74) Agent: WEST, Vivien; SmithKline Beecham, Corporate Intellectual Property, Two New Horizons Court, Brentford, Middlesex TW8 9EP (GB).

## (54) Title: PROCESS FOR THE PREPARATION OF PAROXETINE HYDROCHLORIDE

#### (57) Abstract

Solid forms of paroxetine hydrochloride (crystalline as well as amorphous) are obtained by precipitation from a supercritical or near-critical fluid such as carbon dioxide. This procedure offers the advantage of a more easily controlled precipitation process than is obtained by previously known methods, and better control of surface morphology, porosity, particle size and distribution.

This procedure offers the advantage of a more easily controlled precipitation process than is obtained by previously known methods, and better control of surface morphology, porosity, particle size and distribution. These factors are important and affect, for example, the rate of dissolution and the performance of the material in secondary pharmaceutical manufacturing.

Paroxetine hydrochloride is prepared for supercritical fluid precipitation by forming a solution in a supercritical fluid such as supercritical carbon dioxide. Other supercritical fluids such as ethane, n-propane, n-butane, and nitrogen oxide, may also be used. Known solid forms of paroxetine hydrochloride form such solutions with some difficulty. Therefore the paroxetine hydrochloride is preferably first dissolved in an auxiliary solvent, for example ethanol, propan-2-ol, or isobutyl alcohol, compatible with the supercritical fluid, and the solution brought into contact with the supercritical fluid to form a suitable solution for precipitation. A suitable solution may be prepared from amorphous paroxetine hydrochloride or a crystalline anhydrate, hydrate, or solvate of paroxetine hydrochloride, or by dissolving the free base and hydrochloric acid in an aqueous, organic or mixed aqueous and organic solvent. Indeed it may be possible to bring the free base and hydrochloric acid together in the supercritical domain where they may react prior to precipitation of paroxetine hydrochloride.

20

25

10

15

In order to achieve successful precipitation of paroxetine hydrochloride from a supercritical fluid by a method that uses an auxiliary solvent, this solvent preferably has an affinity for the supercritical fluid so that both may be effectively removed in a single process. The preferred auxiliary solvents listed above are not ideal in this respect, at least when the supercritical fluid is carbon dioxide, so it may be advantageous to employ an additional entraining solvent to confer suitable properties on the auxiliary solvent. An example of a suitable entraining solvent for use with propan-2-ol and supercritical carbon dioxide is acetone.

The entraining solvent may be combined with the auxiliary solvent in a ratio from 1:5 to 20:1, preferably from 1:1 to 10:1, and most preferably from 3:1 to 7:1. The concentration of paroxetine hydrochloride in the auxiliary solvent may be from 0.5% to 25%, but is preferably in the range 1% to 10%, for example from 2.5% to 5%. The supercritical solution for precipitation is formed by combining the paroxetine hydrochloride solution with supercritical fluid in its liquid phase in a ratio of from 1:2 to 1:200, preferably in the range from 1:10 to 1:50, most preferably in the range 1:15 to 1:30.

In a typical procedure, a chamber containing a spray device is maintained at a temperature and pressure such that carbon dioxide (or other fluid) is supercritical. The temperature is controlled using an oven, and the pressure is controlled using a back pressure regulator at the chamber exit. A solution of paroxetine hydrochloride is prepared in a suitable solvent system and this solution and a supercritical fluid are separately metered to the spray device using high pressure pumps. Within, or close to the spray device, the supercritical fluid effectively removes the solvent from the paroxetine hydrochloride solution, giving a precipitate which is deposited in the collection chamber. When sufficient material has accumulated in the collection chamber the delivery of paroxetine hydrochloride solution is stopped. The paroxetine hydrochloride particles are rinsed with supercritical fluid to remove final traces of solvent and the apparatus is depressurised to harvest the product.

10

15

30

35

In an alternative method of operation the paroxetine hydrochloride is dissolved in the supercritical fluid or solvent-modified supercritical fluid using a saturator chamber. The resultant supercritical fluid solution is sprayed through a spray device into a second chamber at atmospheric or slightly above atmospheric pressure and particles of paroxetine hydrochloride are formed.

The temperature of the precipitation is generally from 15°C to 150°C, preferably from 45°C to 100°C, and the pressure is maintained in the range 25 to 300 bar, preferably from 100 to 200 bar.

When a crystalline product is desired, improved control of the precipitation process may be achieved by the addition of seeds. When the desired product is a hydrate such as paroxetine hydrochloride hemihydrate, an amount of water should be present in excess of the amount required according to theory.

The solid product of this invention may be formulated for therapy in the dosage forms described in EP-A-0223403 or WO96/24595. Free-flowing solids are advantageous for the preparation of solid formulations. Easily soluble solids are suitable for the preparation of solutions for parenteral use.

Therapeutic uses of the paroxetine product of this invention include treatment of: alcoholism, anxiety, depression, obsessive compulsive disorder, panic disorder, chronic pain, obesity, senile dementia, migraine, bulimia, anorexia, social phobia, pre-menstrual syndrome (PMS), adolescent depression, trichotillomania, dysthymia, and substance abuse, referred to below as "the disorders".

15

Accordingly, the present invention also provides:

- a pharmaceutical composition for treatment or prophylaxis of the disorders comprising solid paroxetine hydrochloride obtained by the process of this invention and a pharmaceutically acceptable carrier or a solution of the obtained solid paroxetine hydrochloride;
- the use of solid paroxetine hydrochloride obtained by the process of this invention to
  manufacture a medicament in solid or liquid form for the treatment or prophylaxis of
  the disorders; and
  - a method of treating the disorders which comprises administering an effective or prophylactic amount of solid paroxetine hydrochloride obtained by the process of this invention, or a solution thereof, to a person suffering from one or more of the disorders.

The invention is illustrated by the following Examples:

### Example 1

A particle collection chamber incorporating a spray device was maintained at a temperature of 45°C and a pressure of 95 bar. A 2% solution of paroxetine hydrochloride in a 50:10 acetone/propan-2-ol mixture containing 1.6% water was metered to the spray device at 0.40 ml/min. Supercritical carbon dioxide was also metered to the spray device at 9.0 ml/min. Paroxetine hydrochloride was deposited as low bulk density white powder.

10

15

### Example 2

A particle collection chamber with a spray device was maintained at 50°C and 125 bar. A 2% solution of paroxetine hydrochloride in 110:10 acetone/propan-2-ol was metered to the spray device at 0.27 ml/min. Supercritical carbon dioxide was also metered to the spray device at 6.0 ml/min. Paroxetine hydrochloride was deposited as a dense white powder.

#### CLAIMS:

- 1. A process for isolating a solid form of paroxetine hydrochloride which comprises precipitating paroxetine hydrochloride from a solution thereof in a supercritical or near-critical fluid
- 2. A process according to claim 1 in which the solid form of paroxetine hydrochloride is crystalline.
- 3. A process according to claim 1 or 2 in which the supercritical fluid is carbon dioxide or ethane, n-propane, n-butane, or nitrogen oxide.
  - 4. A process according to claim 1, 2 or 3 in which the paroxetine hydrochloride is first dissolved in an auxiliary solvent compatible with the supercritical fluid, and the solution brought into contact with the supercritical fluid to form a suitable solution for precipitation.
    - 5. A process according to claim 4 in which the auxiliary solvent is ethanol, propan-2-ol, or isobutyl alcohol.
    - 6. A process according to claim 4 or 5 in which an additional entraining solvent is employed to confer suitable properties on the auxiliary solvent.
- 7. A process according to claim 6 in which the supercritical fluid is carbon dioxide, the auxiliary solvent is propan-2-ol, and the entraining solvent is acetone.
  - 8. A pharmaceutical composition for treatment or prophylaxis of the disorders comprising solid paroxetine hydrochloride obtained by the process of this invention and a pharmaceutically acceptable carrier or a solution of the obtained solid paroxetine hydrochloride.
  - 9. The use of solid paroxetine hydrochloride obtained by the process of this invention to manufacture a medicament in solid or liquid form for the treatment or prophylaxis of the disorders.
  - 10. A method of treating the disorders which comprises administering an effective or prophylactic amount of solid paroxetine hydrochloride obtained by the process of this invention, or a solution thereof, to a person suffering from one or more of the disorders.

30

20

Inter anal Application No PCT/GB 99/03664

		ј РСТ	/GB 99	/03664		
A. CLASS	FICATION OF SUBJECT MATTER C07D405/12 A61K31/4525					
According to	o International Patent Classification (IPC) or to both national classific	ation and IPC				
	SEARCHED	adori and IPO				
Minimum do IPC 7	cumentation searched (classification system followed by classificati CO7D A61K	on symbols)		- 1930		
	tion searched other than minimum documentation to the extent that $\dot{ ext{s}}$					
Electronic d	ata base consulted during the international search (name of data ba	se and, where practical, search	terms used)			
				•		
C. DOCUM	ENTS CONSIDERED TO BE RELEVANT					
Category *	Citation of document, with indication, where appropriate, of the rele	evant passages		Relevant to claim No.		
Α	WO 96 24595 A (SMITHKLINE BEECHAM ;JACEWICZ VICTOR WITOLD (GB); WAR		1-10			
	(GB) 15 August 1996 (1996-08-15) cited in the application page 6, line 5 - line 10; claims		-			
, <b>A</b>	WO 98 31365 A (WARD NEAL ;JACEWIC WITOLD (GB); SMITHKLINE BEECHAM P 23 July 1998 (1998-07-23)	Z VICTOR LC (GB)		1-10		
<b>A</b>	EP 0 223 403 A (BEECHAM GROUP PLC 27 May 1987 (1987-05-27) cited in the application examples			1,8-10		
	<del></del>	·/				
X Furti	ner documents are listed in the continuation of box C.	X Patent family members	s are listed in	n annex.		
* Special ca	tegories of cited documents :	"T" later document published at	ter the inter	national filing data		
"E" earlier of filing of "L" docume which	ant which may throw doubts on priority claim(s) or	or priority date and not in c cited to understand the prin invention  "X" document of particular relev cannot be considered nove involve an inventive step w  "Y" document of particular relev cannot be considered to in	conflict with the conflict with the conflict with the conflict the conflict wants; the conflict with the conflict wants; the conflict with the conflict wants; the conflict with the conflict wi	he application but ory underlying the simed invention be considered to ument its taken alone aimed invention		
other	ent referring to an oral disclosure, use. exhibition or means are the published prior to the international filing date but hand the priority date claimed	document is combined with ments, such combination be in the art.  18. document member of the sa	h one or mor	e other súch docu- s to a person skilled		
Date of the	actual completion of the international search	Date of malting of the interr	national sea	rch report		
1	3 January 2000	24/01/2000				
Name and I	nailing address of the ISA  European Patent Office, P.B. 5818 Patentiaan 2  NL - 2280 HV Rijswijk  Tel. (+31-70) 340-2040. Tx. 31 651 epo nl,  Fax: (+31-70) 340-3016	Authorized officer  De Jong, B				

1.

Inter anal Application No PCT/GB 99/03664

	INION) DOCUMENTS CONSIDERED TO BE RELEVANT	····
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	EP 0 810 224 A (ASAHI GLASS CO LTD) 3 December 1997 (1997-12-03) cited in the application claim 1	1,8-10
A	WO 95 01221 A (UNIV BRADFORD ; HANNA MAZEN (GB); YORK PETER (GB)) 12 January 1995 (1995-01-12) abstract	1 .
:		
·		
-		
,		

In ...national application No.

PCT/GB 99/03664

Box I	Observations where certain claims were found unsearchable (Continuation of Item 1 of first sheet)
This Inte	ernational Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:
1. X	Claims Nos.:  10 because they relate to subject matter not required to be searched by this Authority, namely:  Remark: Although claim 10  is directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged
2.	effects of the compounds.  Claims Nos.: because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
з. 🗌	Claims Nos.: because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).
Box II	Observations where unity of invention is lacking (Continuation of item 2 of first sheet)
This Inte	ornational Searching Authority found multiple inventions in this international application, as follows:
1.	As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2.	As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3.	As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
• • •	
4.	No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims: it is covered by claims Nos.:
Remark	The additional search fees were accompanied by the applicant's protest.
	No protest accompanied the payment of additional search fees.

information on patent family members

Inter anal Application No PCT/GB 99/03664

	atent document		Publication		Patent family	99/03664 Publication
	in search repor		date		member(s)	date
WO	9624595	Α	15-08-1996	AU	701518 B	28-01-1999
				AU	4332896 A	15-08-1996
	•			AU	4786496 A	27-08-1996
	•			AU	9821398 A	04-03 <b>-</b> 1999
				BE	1009112 A	05-11-1996
	•			BG	100333 A	30-08-1996
				BR	9600534 A	13-05-1997
				CA	2168829 A,C	07-08-1996.
				CA	2210022 A	07-08-1996
				CA	2210023 A,C	07-08-1996
				CA	2211521 A	07-08-1996
				CA	2211522 A	07-08-1996
				CH	688353 A	15-08-1997
		•		CH	689229 A	31-12-1998
			•	CN	1143643 A	26-02-1997
				CY	2015 A	20-02-1998
				CZ	9600320 A	14-08-1996
				DE	19603797 A	14-08-1996
	-			DE	29623383 U	20-05-1998
				DK	11996 A	07-08-1996
			•	EP	0808314 A	26-11-1997
				ES	2114471 A	16-05-1998
				FI	960519 A	07-08-1996
				FR	2730232 A	09-08-1996
				GB	2297550 A,B	07-08-1996
			•	GR	1002466 B	06-11-1996
				HK	59397 A	16-05-1997
				HÙ	9600255 A	28-03-1997
				IE	960104 A	07-08-1996
				ΪŢ	MI960203 A	05-08-1997
				ĴP	2915338 B	05-05-1997
				JP	8245620 A	24-09-1996
				JP	11228571 A	
				LT	96007 A,B	24-08-1999 25-10-1996
•				LU		
					88711 A	23-08-1996
	•			LV	11618 A	20-12-1996
				LV	11618 B	20-04-1997
			•	MC	2411 A	02-12-1996
			•	NL	1002248 C	11-09-1996
				NL	1002248 A	06-08-1996
				NO	960472 A	07-08-1996
				NZ	280943 A	29-01-1997
				PL	312646 A	19-08-1996
				PT	101827 A,B	30-09-1996
				RO	112426 A	30-09-1997
			•	SE	9600406 A	07-08-1996
				SG	43787 A	14-11-1997
				SI 	9600036 A	31-10-1996
WO	9831365	Α	23-07-1998	AU	5567398 A	07-08-1998
				EP	0952831 A	03-11-1999
				NO	993460 A	14-09-1999
ΕP	0223403	Α	27-05-1987	AU	593295 B	08-02-1990
				AU	6433286 A	30-04-1987
				BG	61323 B	30-05-1997
				CA	1287060 A	30-07-1991
				ĊZ	9103910 A	19-01-1994
				~ ~~	· · · · · · · · · · · · · · · · ·	

information on patent family members

Inter onal Application No
PCT/GB 99/03664

		<del></del>			99/03004	
Patent document cited in search report		Publication date		t family ber(s)	Publication date	
EP 0223403	A		DE 3 DK DK ES 2 FI HK IE JP 1 JP 6 JP 62 NO NZ PT	1743 A 688827 A 688827 T 61091 A 508786 A 058061 T 864320 A,B, 125993 A 59901 B 918281 C 047587 B 129280 A 864237 A,B, 218047 A 83608 A,B 721723 A	17-02-1995 09-09-1993 31-03-1994 05-04-1991 26-04-1987 01-11-1994 26-04-1987 19-11-1993 20-04-1994 07-04-1995 22-06-1994 11-06-1987 27-04-1987 29-03-1989 01-11-1986 26-01-1988	
EP 0810224	A	03-12-1997		 206592 A 045756 A	30-11-1997 17-02-1998	
WO 9501221	A	12-01-1995	AU 7 CA 2 DE 69 DE 69 EP 0 ES 2 GR 3 JP 8 NZ SG	174530 T 677345 B 007194 A 166301 A 415320 D 415320 T 706421 A 128564 T 029612 T 511987 T 267697 A 47392 A 851453 A	15-01-1999 17-04-1997 24-01-1995 12-01-1995 28-01-1999 24-06-1999 17-04-1996 16-05-1999 30-06-1999 17-12-1996 26-05-1997 17-04-1998 22-12-1998	